# antibodies -online.com





# anti-DDX42 antibody (C-Term)

2 Images



Go to Product page

$\sim$						
	1//	Д	r۱	/1	$\triangle$	٨

**Target Details** 

DDX42

Target:


Quantity: 100 µL Target: DDX42  Binding Specificity: C-Term  Reactivity: Human, Mouse  Host: Rabbit Clonality: Polyclonal  Conjugate: This DDX42 antibody is un-conjugated  Application: Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC), Immunocytochemistry (ICC)  Product Details  Immunogen: A synthesized peptide derived from human DDX42, corresponding to a region within C-termina amino acids.  Isotype: IgG  Specificity: DDX42 Antibody detects endogenous levels of total DDX42.  Predicted Reactivity: Pig,Bovine,Horse,Rabbit,Dog  Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).			
Binding Specificity: C-Term  Reactivity: Human, Mouse  Host: Rabbit  Clonality: Polyclonal  Conjugate: This DDX42 antibody is un-conjugated  Application: Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC), Immunocytochemistry (ICC)  Product Details  Immunogen: A synthesized peptide derived from human DDX42, corresponding to a region within C-termina amino acids.  Isotype: IgG  Specificity: DDX42 Antibody detects endogenous levels of total DDX42.  Predicted Reactivity: Pig,Bovine,Horse,Rabbit,Dog  Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling	Quantity:	100 μL	
Reactivity: Human, Mouse  Host: Rabbit  Clonality: Polyclonal  Conjugate: This DDX42 antibody is un-conjugated  Application: Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC), Immunocytochemistry (ICC)  Product Details  Immunogen: A synthesized peptide derived from human DDX42, corresponding to a region within C-termina amino acids.  Isotype: IgG  Specificity: DDX42 Antibody detects endogenous levels of total DDX42.  Predicted Reactivity: Pig,Bovine,Horse,Rabbit,Dog  Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling	Target:	DDX42	
Host: Rabbit  Clonality: Polyclonal  Conjugate: This DDX42 antibody is un-conjugated  Application: Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC), Immunocytochemistry (ICC)  Product Details  Immunogen: A synthesized peptide derived from human DDX42, corresponding to a region within C-termina amino acids.  Isotype: IgG  Specificity: DDX42 Antibody detects endogenous levels of total DDX42.  Predicted Reactivity: Pig,Bovine,Horse,Rabbit,Dog  Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling	Binding Specificity:	C-Term	
Clonality: Polyclonal  Conjugate: This DDX42 antibody is un-conjugated  Application: Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC), Immunocytochemistry (ICC)  Product Details  Immunogen: A synthesized peptide derived from human DDX42, corresponding to a region within C-termina amino acids.  Isotype: IgG  Specificity: DDX42 Antibody detects endogenous levels of total DDX42.  Predicted Reactivity: Pig,Bovine,Horse,Rabbit,Dog  Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling	Reactivity:	Human, Mouse	
Conjugate: This DDX42 antibody is un-conjugated  Application: Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC), Immunocytochemistry (ICC)  Product Details  Immunogen: A synthesized peptide derived from human DDX42, corresponding to a region within C-termina amino acids.  Isotype: IgG  Specificity: DDX42 Antibody detects endogenous levels of total DDX42.  Predicted Reactivity: Pig,Bovine,Horse,Rabbit,Dog  Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling	Host:	Rabbit	
Application:  Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC), Immunocytochemistry (ICC)  Product Details  Immunogen:  A synthesized peptide derived from human DDX42, corresponding to a region within C-termina amino acids.  Isotype:  IgG  Specificity:  DDX42 Antibody detects endogenous levels of total DDX42.  Predicted Reactivity:  Pig,Bovine,Horse,Rabbit,Dog  Purification:  The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling	Clonality:	Polyclonal	
Immunocytochemistry (ICC)  Product Details  Immunogen:  A synthesized peptide derived from human DDX42, corresponding to a region within C-termina amino acids.  Isotype:  IgG  Specificity:  DDX42 Antibody detects endogenous levels of total DDX42.  Predicted Reactivity:  Pig,Bovine,Horse,Rabbit,Dog  Purification:  The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling	Conjugate:	This DDX42 antibody is un-conjugated	
Immunogen:  A synthesized peptide derived from human DDX42, corresponding to a region within C-termina amino acids.  Isotype:  IgG  Specificity:  DDX42 Antibody detects endogenous levels of total DDX42.  Predicted Reactivity:  Pig,Bovine,Horse,Rabbit,Dog  Purification:  The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling	Application:		
amino acids.  Isotype: IgG  Specificity: DDX42 Antibody detects endogenous levels of total DDX42.  Predicted Reactivity: Pig,Bovine,Horse,Rabbit,Dog  Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling	Product Details		
Specificity: DDX42 Antibody detects endogenous levels of total DDX42.  Predicted Reactivity: Pig,Bovine,Horse,Rabbit,Dog  Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling	Immunogen:	A synthesized peptide derived from human DDX42, corresponding to a region within C-terminal amino acids.	
Predicted Reactivity: Pig,Bovine,Horse,Rabbit,Dog  Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling	Isotype:	IgG	
Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling	Specificity:	DDX42 Antibody detects endogenous levels of total DDX42.	
	Predicted Reactivity:	Pig,Bovine,Horse,Rabbit,Dog	
	Purification:		

## **Target Details**

Expiry Date:


Target Details				
Alternative Name:	DDX42 (DDX42 Products)			
Background:	Description: ATP-dependent RNA helicase. Binds to partially double-stranded RNAs (dsRNAs) ir			
	order to unwind RNA secondary structures. Unwinding is promoted in the presence of single-			
	strand binding proteins. Mediates also RNA duplex formation thereby displacing the single-			
	strand RNA binding protein. ATP and ADP modulate its activity: ATP binding and hydrolysis by			
	DDX42 triggers RNA strand separation, whereas the ADP-bound form of the protein triggers			
	annealing of complementary RNA strands. Involved in the survival of cells by interacting with			
	TP53BP2 and thereby counteracting the apoptosis-stimulating activity of TP53BP2. Relocalizes			
	TP53BP2 to the cytoplasm.			
	Gene: DDX42			
Molecular Weight:	103 kDa			
Gene ID:	11325			
UniProt:	Q86XP3			
Application Details				
Application Notes:	WB 1:1000-3000, IHC 1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000			
Restrictions:	For Research Use only			
Handling				
Format:	Liquid			
Concentration:	1 mg/mL			
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %			
	glycerol.			
Preservative:	Sodium azide			
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which			
	should be handled by trained staff only.			
Storage:	-20 °C			
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.			

12 months



### **Immunohistochemistry**

**Image 1.** ABIN6279173 at 1/100 staining Mouse muscle tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at  $22_{\rm i}$  aC. An HRP conjugated goat anti-rabbit antibody was used as the secondary



#### **Western Blotting**

**Image 2.** Western blot analysis of DDX42 using HuvEc whole cell lysates