# antibodies .- online.com







## anti-ZDHHC9 antibody (C-Term)

**Images** 

Overview

Target Details

ZDHHC9

Target:



Quantity:	100 μL
Target:	ZDHHC9
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZDHHC9 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized peptide derived from human DHHC-9, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	DHHC-9 Antibody detects endogenous levels of total DHHC-9.
Predicted Reactivity:	Pig,Bovine,Horse,Rabbit
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

#### **Target Details**


Alternative Name:	ZDHHC9 (ZDHHC9 Products)
Background:	Description: The ZDHHC9-GOLGA7 complex is a palmitoyltransferase specific for HRAS and NRAS.  Gene: ZDHHC9
Molecular Weight:	41kDa
Gene ID:	51114
UniProt:	Q9Y397

### Application Details

Application Notes:	WB 1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

### Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



# kDa 250— 150— 100— 75— 50— 37—

15-

#### Immunofluorescence (fixed cells)

**Image 1.** ABIN6272287 staining COLO205 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

#### **Western Blotting**

**Image 2.** Western blot analysis of DHHC-9 using COLO205 whole cell lysates