antibodies -online.com





anti-DLGAP5 antibody (N-Term)

3 Images



Go to Product page

Overview

Quantity:	100 μL
Target:	DLGAP5
Binding Specificity:	N-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DLGAP5 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), ELISA, Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human DLGAP5, corresponding to a region within N-terminal amino acids.
Isotype:	IgG
Specificity:	DLGAP5 Antibody detects endogenous levels of total DLGAP5.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog,Chicken
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

- .	DI CARE	
Target:	DLGAP5	

Target Details


Alternative Name:	DLGAP5 (DLGAP5 Products)
Background:	Description: Potential cell cycle regulator that may play a role in carcinogenesis of cancer cells.
	Mitotic phosphoprotein regulated by the ubiquitin-proteasome pathway. Key regulator of
	adherens junction integrity and differentiation that may be involved in CDH1-mediated adhesion
	and signaling in epithelial cells.
	Gene: DLGAP5
Molecular Weight:	95kDa
Gene ID:	9787
UniProt:	Q15398
Pathways:	M Phase

Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months




Western Blotting

Image 1. Western blot analysis of extracts of HEK-293, using DLGAP5 antibody. The lane on the left is treated with the antigen-specific peptide.



Immunofluorescence (fixed cells)

Image 2. ABIN6277179 staining Hela cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibod



Immunohistochemistry

Image 3. ABIN6277179 at 1/100 staining Mouse liver tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22_iaC. An HRP conjugated goat anti-rabbit antibody was used as the secondary