# antibodies .- online.com







## anti-Dystonin antibody (C-Term)

**Images** 



_		
()Ver	view	

Overview	
Quantity:	100 μL
Target:	Dystonin (DST)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Dystonin antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized peptide derived from human DST, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	DST Antibody detects endogenous levels of total DST.
Predicted Reactivity:	Pig,Bovine,Horse,Dog,Chicken

## **Target Details**

Purification:


Target: Dystonin (DST)

Resin (Thermo Fisher Scientific).

The antiserum was purified by peptide affinity chromatography using SulfoLink<sup>TM</sup> Coupling


## **Target Details**

Alternative Name:	DST (DST Products)
Background:	Description: Cytoskeletal linker protein. Acts as an integrator of intermediate filaments, actin
	and microtubule cytoskeleton networks. Required for anchoring either intermediate filaments to
	the actin cytoskeleton in neural and muscle cells or keratin-containing intermediate filaments to
	hemidesmosomes in epithelial cells. The proteins may self-aggregate to form filaments or a
	two-dimensional mesh. Regulates the organization and stability of the microtubule network of
	sensory neurons to allow axonal transport. Mediates docking of the dynein/dynactin motor
	complex to vesicle cargos for retrograde axonal transport through its interaction with
	TMEM108 and DCTN1 (By similarity).
	Gene: DST
Molecular Weight:	861kDa
Gene ID:	667
UniProt:	Q03001
Application Details	
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %
	glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



### **Western Blotting**

**Image 1.** Western blot analysis of extracts from A431, using DST Antibody.



#### **Immunohistochemistry**

**Image 2.** ABIN6277015 at 1/100 staining Human lung tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22<sub>i</sub>aC. An HRP conjugated goat anti-rabbit antibody was used as the secondary