antibodies -online.com









Overview Quantity:

Quantity:	100 μL
Target:	EHD1
Binding Specificity:	C-Term
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EHD1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human EHD1, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	EHD1 Antibody detects endogenous levels of total EHD1.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

|--|


Target Details

Alternative Name:	EHD1 (EHD1 Products)
Background:	Description: ATP- and membrane-binding protein that controls membrane
	reorganization/tubulation upon ATP hydrolysis. In vitro causes vesiculation of endocytic
	membranes (PubMed:24019528). Acts in early endocytic membrane fusion and membrane
	trafficking of recycling endosomes (PubMed:15020713, PubMed:17233914,
	PubMed:20801876). Recruited to endosomal membranes upon nerve growth factor
	stimulation, indirectly regulates neurite outgrowth (By similarity). Plays a role in myoblast fusion
	(By similarity). Involved in the unidirectional retrograde dendritic transport of endocytosed
	BACE1 and in efficient sorting of BACE1 to axons implicating a function in neuronal APP
	processing (By similarity). Plays a role in the formation of the ciliary vesicle (CV), an early step in
	cilium biogenesis. Proposed to be required for the fusion of distal appendage vesicles (DAVs) to
	form the CV by recruiting SNARE complex component SNAP29. Is required for recruitment of
	transition zone proteins CEP290, RPGRIP1L, TMEM67 and B9D2, and of IFT20 following DAV
	reorganization before Rab8-dependent ciliary membrane extension. Required for the loss of
	CCP110 form the mother centriole essential for the maturation of the basal body during
	ciliogenesis (PubMed:25686250).
	Gene: EHD1
Molecular Weight:	61 kDa
Gene ID:	10938
UniProt:	Q9H4M9
Pathways:	Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development
Application Details	
Application Notes:	WB 1:1000-3000, IHC 1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %
	glycerol.

Handling


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. ABIN6278061 at 1/100 staining RAT testis tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at $22_i\tilde{a}C$. An HRP conjugated goat anti-rabbit antibody was used as the secondary



Western Blotting

Image 2. Western blot analysis of EHD1 using RAW264.7 whole lysates