# ANTIBODIES ONLINE

# Datasheet for ABIN6261631 anti-EZH2 antibody (Internal Region)

2 Images



Overview

Quantity:	100 µL
Target:	EZH2
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EZH2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

# Product Details

Immunogen:	A synthesized peptide derived from human EZH2, corresponding to a region within the internal amino acids.
Isotype:	lgG
Specificity:	EZH2 Antibody detects endogenous levels of total EZH2.
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Rabbit,Dog,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

Target Details

Target: EZH2

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6261631 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Alternative Name:	EZH2 (EZH2 Products)
Background:	Description: Polycomb group (PcG) protein. Catalytic subunit of the PRC2/EED-EZH2 complex,
	which methylates 'Lys-9' (H3K9me) and 'Lys-27' (H3K27me) of histone H3, leading to
	transcriptional repression of the affected target gene. Able to mono-, di- and trimethylate 'Lys-
	27' of histone H3 to form H3K27me1, H3K27me2 and H3K27me3, respectively. Displays a
	preference for substrates with less methylation, loses activity when progressively more methyl
	groups are incorporated into H3K27, H3K27me0 > H3K27me1 > H3K27me2
	(PubMed:22323599). Compared to EZH1-containing complexes, it is more abundant in
	embryonic stem cells and plays a major role in forming H3K27me3, which is required for
	embryonic stem cell identity and proper differentiation. The PRC2/EED-EZH2 complex may also
	serve as a recruiting platform for DNA methyltransferases, thereby linking two epigenetic
	repression systems. Genes repressed by the PRC2/EED-EZH2 complex include HOXC8, HOXA9,
	MYT1, CDKN2A and retinoic acid target genes. EZH2 can also methylate non-histone proteins
	such as the transcription factor GATA4 and the nuclear receptor RORA. Regulates the circadian
	clock via histone methylation at the promoter of the circadian genes. Essential for the CRY1/2-
	mediated repression of the transcriptional activation of PER1/2 by the CLOCK-ARNTL/BMAL1
	heterodimer, involved in the di and trimethylation of 'Lys-27' of histone H3 on PER1/2
	promoters which is necessary for the CRY1/2 proteins to inhibit transcription.
	Gene: EZH2
Molecular Weight:	85 kDa
Gene ID:	2146
UniProt:	Q15910
Pathways:	Retinoic Acid Receptor Signaling Pathway, Regulation of Muscle Cell Differentiation
Application Details	
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN6261631 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

## Handling

	glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

### Images



### Immunohistochemistry

**Image 1.** ABIN6268696 at 1/100 staining Mouse testis tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



#### Western Blotting

**Image 2.** Western blot analysis of EZH2 expression in HEK293 cell lysate,The lane on the left is treated with the antigen-specific peptide.