antibodies.com

Datasheet for ABIN6261731 anti-FGFR1 antibody (C-Term)

5 Images



Overview

Quantity:	100 µL		
Target:	FGFR1		
Binding Specificity:	C-Term		
Reactivity:	Human, Mouse, Rat		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This FGFR1 antibody is un-conjugated		
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)		

Product Details

Immunogen:	A synthesized peptide derived from human FGFR1, corresponding to a region within C-terminal amino acids.
Isotype:	lgG
Specificity:	FGFR1 Antibody detects endogenous levels of total FGFR1.
Predicted Reactivity:	Bovine,Horse,Sheep,Rabbit,Dog,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

To	-	~+·
1.2	41(1	ω
	. 9	οι.

FGFR1

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN6261731 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
Alternative Name:	FGFR1 (FGFR1 Products)
Background:	Description: Tyrosine-protein kinase that acts as cell-surface receptor for fibroblast growth
	factors and plays an essential role in the regulation of embryonic development, cell
	proliferation, differentiation and migration. Required for normal mesoderm patterning and
	correct axial organization during embryonic development, normal skeletogenesis and normal
	development of the gonadotropin-releasing hormone (GnRH) neuronal system. Phosphorylates
	PLCG1, FRS2, GAB1 and SHB. Ligand binding leads to the activation of several signaling
	cascades. Activation of PLCG1 leads to the production of the cellular signaling molecules
	diacylglycerol and inositol 1,4,5-trisphosphate. Phosphorylation of FRS2 triggers recruitment of
	GRB2, GAB1, PIK3R1 and SOS1, and mediates activation of RAS, MAPK1/ERK2, MAPK3/ERK1
	and the MAP kinase signaling pathway, as well as of the AKT1 signaling pathway. Promotes
	phosphorylation of SHC1, STAT1 and PTPN11/SHP2. In the nucleus, enhances RPS6KA1 and
	CREB1 activity and contributes to the regulation of transcription. FGFR1 signaling is down-
	regulated by IL17RD/SEF, and by FGFR1 ubiquitination, internalization and degradation.
	Gene: FGFR1
Molecular Weight:	120kDa
Gene ID:	2260
UniProt:	P11362
Pathways:	RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin
	Signaling Pathway, Sensory Perception of Sound, Stem Cell Maintenance, S100 Proteins

Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

Handling


Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN6261731 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

lond	lina
land	
10110	


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. ABIN6269136 at 1/100 staining Rat kidney tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



Immunofluorescence (fixed cells)

Image 2. ABIN6269136 staining A549 cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25jãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37jãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibod

Images

kDa	1	2	
250-			
150-		-	
100-			
75-			
50-			
37—			
25-			
20-			
15-			

Western Blotting

Image 3. Western blot analysis of FGFR1 expression in HepG2 whole cell lysates,The lane on the left is treated with the antigen-specific peptide.

Please check the product details page for more images. Overall 5 images are available for ABIN6261731.