antibodies - online.com







anti-FOXP1 antibody (C-Term)





Go to	D		
	Pron	וויאווו	nane

()	11	\sim	rv		۱ ۸
	1 \ /	┙	I \/	╙	1/1

Quantity:	100 μL
Target:	FOXP1
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FOXP1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human FOXP1, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	FOXP1 Antibody detects endogenous levels of total FOXP1.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Dog,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	FOXP1	

Alternative Name:	FOXP1 (FOXP1 Products)
Background:	Description: Transcriptional repressor (PubMed:18347093, PubMed:26647308). Can act with
	CTBP1 to synergistically repress transcription but CTPBP1 is not essential (By similarity). Plays
	an important role in the specification and differentiation of lung epithelium. Acts cooperatively
	with FOXP4 to regulate lung secretory epithelial cell fate and regeneration by restricting the
	goblet cell lineage program, the function may involve regulation of AGR2. Essential
	transcriptional regulator of B-cell development. Involved in regulation of cardiac muscle cell
	proliferation. Involved in the columnar organization of spinal motor neurons. Promotes the
	formation of the lateral motor neuron column (LMC) and the preganglionic motor column (PGC)
	and is required for respective appropriate motor axon projections. The segment-appropriate
	generation of spinal chord motor columns requires cooperation with other Hox proteins. Can
	regulate PITX3 promoter activity, may promote midbrain identity in embryonic stem cell-derived
	dopamine neurons by regulating PITX3. Negatively regulates the differentiation of T follicular
	helper cells T(FH)s. Involved in maintenance of hair follicle stem cell quiescence, the function
	probably involves regulation of FGF18 (By similarity). Represses transcription of various pro-
	apoptotic genes and cooperates with NF-kappa B-signaling in promoting B-cell expansion by
	inhibition of caspase-dependent apoptosis (PubMed:25267198). Binds to CSF1R promoter
	elements and is involved in regulation of monocyte differentiation and macrophage functions,
	repression of CSF1R in monocytes seems to involve NCOR2 as corepressor
	(PubMed:15286807, PubMed:18799727, PubMed:18347093). Involved in endothelial cell
	proliferation, tube formation and migration indicative for a role in angiogenesis, the role in
	neovascularization seems to implicate suppression of SEMA5B (PubMed:24023716). Can
	negatively regulate androgen receptor signaling (PubMed:18640093).
	Gene: FOXP1
Molecular Weight:	75kDa
Gene ID:	27086
UniProt:	Q9H334
Pathways:	Chromatin Binding, Regulation of Muscle Cell Differentiation, Positive Regulation of Immune
	Effector Process, Production of Molecular Mediator of Immune Response
Application Details	
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

Images




100 — Hat braits 100 — Hat braits 50 — Peptide

Immunofluorescence (fixed cells)

Image 1. ABIN6277494 staining Hela cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600 was used as secondary antibod

Western Blotting

Image 2. Western blot analysis of extracts from rat brain, using FOXP1 Antibody.



Immunohistochemistry

Image 3. ABIN6277494 at 1/100 staining Mouse spleen tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22¡ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary

Please check the product details page for more images. Overall 4 images are available for ABIN6261807.