



[Go to Product page](#)

Datasheet for ABIN6261824
anti-FSTL1 antibody (N-Term)

2 Images

Overview

Quantity:	100 µL
Target:	FSTL1
Binding Specificity:	N-Term
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FSTL1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Immunogen:	A synthesized peptide derived from human FSTL1, corresponding to a region within N-terminal amino acids.
Isotype:	IgG
Specificity:	FSTL1 Antibody detects endogenous levels of total FSTL1.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	FSTL1
---------	-------

Target Details


Alternative Name:	FSTL1 (FSTL1 Products)
Background:	Description: May modulate the action of some growth factors on cell proliferation and differentiation. Binds heparin (By similarity). Gene: FSTL1
Molecular Weight:	35-37 kDa
Gene ID:	11167
UniProt:	Q12841

Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



Western Blotting

Image 1. Western blot analysis of extracts of A2780 cells, using FSTL1 antibody.



Immunohistochemistry

Image 2. ABIN6273118 at 1/100 staining Human Melanoma tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.