
antibodies .- online.com







anti-GAD antibody (C-Term)





\sim	
()\/△	rview
\cup	1 410 44

Target:

Quantity:	100 μL
Target:	GAD (GAD1)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GAD antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF)
Product Details	
Immunogen:	A synthesized peptide derived from human GAD1, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	GAD1 Antibody detects endogenous levels of total GAD1.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).
Target Details	

GAD (GAD1)

Target Details


Alternative Name:	GAD1 (GAD1 Products)
Background:	Description: Catalyzes the production of GABA. Gene: GAD1
Molecular Weight:	67kDa
Gene ID:	2571
UniProt:	Q99259

Application Details

Application Notes:	WB: 1:500-1:3000, IHC: 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months




Immunofluorescence (fixed cells)

Image 1. ABIN6266543 staining LOVO by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



Immunohistochemistry

Image 2. ABIN6266543 at 1/100 staining human brain tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



Western Blotting

Image 3. Western blot analysis of extracts from mouse liver, using GAD1 Antibody.

Please check the product details page for more images. Overall 4 images are available for ABIN6261864.