

Datasheet for ABIN6262000

anti-Glycogen Synthase 1 antibody (C-Term)**3** Images[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Glycogen Synthase 1 (GYS1)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Glycogen Synthase 1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human Glycogen Synthase, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	Glycogen Synthase Antibody detects endogenous levels of total Glycogen Synthase.
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Sheep,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	Glycogen Synthase 1 (GYS1)
---------	----------------------------

Target Details


Alternative Name:	GYS1 (GYS1 Products)
Background:	Description: Transfers the glycosyl residue from UDP-Glc to the non-reducing end of alpha-1,4-glucan. Gene: GYS1
Molecular Weight:	85,89kDa
Gene ID:	2997
UniProt:	P13807
Pathways:	PI3K-Akt Signaling , AMPK Signaling , Cellular Glucan Metabolic Process

Application Details

Application Notes:	WB 1:500-1:2000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months




Western Blotting

Image 1. Western blot analysis of Glycogen Synthase expression in Serum treated HeLa whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



Immunofluorescence (fixed cells)

Image 2. ABIN6269143 staining NIH-3T3 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



Western Blotting

Image 3. Western blot analysis of extracts from various samples, using Glycogen Synthase Antibody. Lane 1: HepG2 treated with blocking peptide; Lane 2: HepG2; Lane 3: B16f10; Lane 4: Mouse brain.