antibodies -online.com





anti-GNG12 antibody

2 Images



Go to Product page

Overview

Quantity:	100 μL
Target:	GNG12
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GNG12 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	A synthesized peptide derived from human GNG12
Isotype:	IgG
Specificity:	GNG12 Antibody detects endogenous levels of total GNG12
Cross-Reactivity:	Human, Mouse (Murine)
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	GNG12
Alternative Name:	GNG12 (GNG12 Products)
Background:	Description: Guanine nucleotide-binding proteins (G proteins) are involved as a modulator or

Target Details


	transducer in various transmembrane signaling systems. The beta and gamma chains are
	required for the GTPase activity, for replacement of GDP by GTP, and for G protein-effector
	interaction.
	Gene: GNG12
Molecular Weight:	8 kDa
Gene ID:	55970
UniProt:	Q9UBI6

Application Details

Application Notes:	WB 1:1000-3000 IHC 1:200
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C.Stable for 12 months from date of receipt
Expiry Date:	12 months



Immunohistochemistry

Image 1. ABIN6279468 at 1/100 staining Human liver cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22¡ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary



Western Blotting

Image 2. Western blot analysis GNG12 using COS7 whole cell lysates