# antibodies .- online.com







## anti-HLF antibody

**Images** 



#### Overview

Quantity:	100 μL
Target:	HLF
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HLF antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

## Product Details

Immunogen:	A synthesized peptide derived from human HLF
Isotype:	IgG
Specificity:	HLF Antibody detects endogenous levels of total HLF.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog,Chicken
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

## **Target Details**

Target:	HLF
Alternative Name:	HLF (HLF Products)
Background:	Gene: HLF

## **Target Details**

Molecular Weight:	33 kDa
Gene ID:	3131
UniProt:	Q16534


## **Application Details**

Application Notes:	WB 1:1000-3000, IF/ICC 1:200-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

### **Images**



### **Western Blotting**

**Image 1.** Western blot analysis of HLF using HeLa whole lysates.



#### Immunofluorescence (fixed cells)

**Image 2.** ABIN6278024 staining HepG2 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod