antibodies .- online.com







anti-IRF3 antibody (C-Term)





\sim						
	1//	Д	r۱	1	Θ 1	٨

Target Details

IRF3

Target:

Quantity:	100 μL
Target:	IRF3
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IRF3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Product Details Immunogen:	A synthesized peptide derived from human IRF3, corresponding to a region within C-terminal amino acids.
Immunogen:	amino acids.
Immunogen: Isotype:	amino acids.


Target Details

Alternative Name:	IRF3 (IRF3 Products)	
Background:	Description: Key transcriptional regulator of type I interferon (IFN)-dependent immune	
	responses which plays a critical role in the innate immune response against DNA and RNA	
	viruses. Regulates the transcription of type I IFN genes (IFN-alpha and IFN-beta) and IFN-	
	stimulated genes (ISG) by binding to an interferon-stimulated response element (ISRE) in their	
	promoters. Acts as a more potent activator of the IFN-beta (IFNB) gene than the IFN-alpha	
	(IFNA) gene and plays a critical role in both the early and late phases of the IFNA/B gene	
	induction. Found in an inactive form in the cytoplasm of uninfected cells and following viral	
	infection, double-stranded RNA (dsRNA), or toll-like receptor (TLR) signaling, is phosphorylated	
	by IKBKE and TBK1 kinases. This induces a conformational change, leading to its dimerization	
	and nuclear localization and association with CREB binding protein (CREBBP) to form dsRNA-	
	activated factor 1 (DRAF1), a complex which activates the transcription of the type I IFN and	
	ISG genes. Can activate distinct gene expression programs in macrophages and can induce	
	significant apoptosis in primary macrophages.	
	Gene: IRF3	
Molecular Weight:	57kDa	
Gene ID:	3661	
UniProt:	Q14653	
Pathways:	TLR Signaling, Activation of Innate immune Response, Cellular Response to Molecule of	
	Bacterial Origin, Hepatitis C, Toll-Like Receptors Cascades	
Application Details		
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %	
	glycerol.	
Preservative:	Sodium azide	

Handling


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.	
Expiry Date:	12 months	

Images




Western Blotting

Image 1. Western blot analysis of IRF3 expression in K562 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



Immunohistochemistry

Image 2. ABIN6269358 at 1/100 staining Human lung tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



Immunofluorescence (fixed cells)

Image 3. ABIN6269358 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibod

Please check the product details page for more images. Overall 4 images are available for ABIN6262651.