

Datasheet for ABIN6262768  
**anti-Keratin 10 antibody (Internal Region)**[Go to Product page](#)

## 3 Images

## Overview

Quantity:	100 µL
Target:	Keratin 10 (KRT10)
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Keratin 10 antibody is un-conjugated
Application:	Immunofluorescence (IF), Immunohistochemistry (IHC), Western Blotting (WB), ELISA, Immunocytochemistry (ICC)

## Product Details

Immunogen:	A synthesized peptide derived from human Cytokeratin 10, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	Cytokeratin 10 Antibody detects endogenous levels of total Cytokeratin 10.
Predicted Reactivity:	Pig,Bovine,Sheep,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

## Target Details

Target:	Keratin 10 (KRT10)
---------	--------------------

## Target Details


Alternative Name:	KRT10 ( <a href="#">KRT10 Products</a> )
Background:	Description: Plays a role in the establishment of the epidermal barrier on plantar skin. Gene: KRT10
Molecular Weight:	59kDa
Gene ID:	3858
UniProt:	<a href="#">P13645</a>

## Application Details

Application Notes:	WB: 1:500-1:3000, IF/ICC: 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months




#### Immunofluorescence (fixed cells)

**Image 1.** ABIN6266577 staining HeLa by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



#### Western Blotting

**Image 2.** Western blot analysis on HeLa cell lysate using Keratin 10 Antibody. The lane on the left is treated with the antigen-specific peptide.



#### Western Blotting

**Image 3.** Western blot analysis of extracts of various tissue, using Keratin 10 antibody.