antibodies .- online.com





anti-LHX8 antibody (Internal Region)

2 Images



Overview 100 μL Quantity: Target: LHX8 Binding Specificity: Internal Region Reactivity: Human, Mouse, Rat Host: Rabbit Clonality: Polyclonal Conjugate: This LHX8 antibody is un-conjugated Application: Western Blotting (WB), ELISA, Immunohistochemistry (IHC) **Product Details** Immunogen: A synthesized peptide derived from human LHX8, corresponding to a region within the internal amino acids. Isotype: IgG Specificity: LHX8 Antibody detects endogenous levels of total LHX8. Predicted Reactivity: Pig, Bovine, Horse, Sheep, Rabbit, Dog, Chicken, Xenopus The antiserum was purified by peptide affinity chromatography using SulfoLinkTM Coupling Purification: Resin (Thermo Fisher Scientific). **Target Details** LHX8 Target:

Target Details


Alternative Name:	LHX8 (LHX8 Products)
Background:	Description: Transcription factor involved in differentiation of certain neurons and mesenchymal cells. Gene: LHX8
Molecular Weight:	43kDa
Gene ID:	431707
UniProt:	Q68G74

Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



Immunohistochemistry

Image 1. ABIN6277041 at 1/100 staining Rat lung tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22¡ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary



Western Blotting

Image 2. Western blot analysis of MCF7 cell lysate, using LHX8 Antibody. The lane on the left is treated with the antigen-specific peptide.