antibodies - online.com







anti-Merlin antibody (C-Term)


Images




Overview	
Quantity:	100 μL
Target:	Merlin (NF2)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Merlin antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized peptide derived from human NF2 / Merlin, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	NF2 / Merlin Antibody detects endogenous levels of total NF2 / Merlin.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	Merlin (NF2)

Target Details

Alternative Name:	NF2 (NF2 Products)
Background:	Description: Probable regulator of the Hippo/SWH (Sav/Wts/Hpo) signaling pathway, a
	signaling pathway that plays a pivotal role in tumor suppression by restricting proliferation and
	promoting apoptosis. Along with WWC1 can synergistically induce the phosphorylation of
	LATS1 and LATS2 and can probably function in the regulation of the Hippo/SWH
	(Sav/Wts/Hpo) signaling pathway. May act as a membrane stabilizing protein. May inhibit PI3
	kinase by binding to AGAP2 and impairing its stimulating activity. Suppresses cell proliferation
	and tumorigenesis by inhibiting the CUL4A-RBX1-DDB1-VprBP/DCAF1 E3 ubiquitin-protein
	ligase complex.
	Gene: NF2
Molecular Weight:	70kDa
Gene ID:	4771
UniProt:	P35240
Pathways:	Cell-Cell Junction Organization
Application Details	
Application Notes:	WB 1:500-1:2000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %
	glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months





Immunofluorescence (fixed cells)

Image 1. ABIN6269222 staining NIH-3T3 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

Western Blotting

Image 2. Western blot analysis of Merlin expression in IFN- α treated HuvEc whole cell lysates,The lane on the left is treated with the antigen-specific peptide.