antibodies.com

Datasheet for ABIN6263352 anti-MRFAP1 antibody (Internal Region)

2 Images



Overview

Quantity:	100 µL
Target:	MRFAP1
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MRFAP1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	A synthesized peptide derived from human MRFAP1, corresponding to a region within the internal amino acids.
lsotype:	lgG
Specificity:	MRFAP1 Antibody detects endogenous levels of total MRFAP1.
Predicted Reactivity:	Bovine,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).


Target Details

Target:

MRFAP1

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6263352 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
Alternative Name:	MRFAP1 (MRFAP1 Products)
Background:	Gene: MRFAP1
Molecular Weight:	15 kDa
Gene ID:	93621
UniProt:	Q9Y605
Application Details	
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



Immunohistochemistry

Image 1. ABIN6273268 at 1/100 staining Mouse lung tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

Western Blotting

Image 2. Western blot analysis of extracts of Jurkat cells, using MRFAP1 antibody.

