# antibodies - online.com







# anti-MVP antibody (N-Term)

**Images** 



Publication



Overview	
Quantity:	100 μL
Target:	MVP
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MVP antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA
Product Details	
Immunogen:	A synthesized peptide derived from human MVP, corresponding to a region within N-terminal amino acids.
Isotype:	IgG
Specificity:	MVP Antibody detects endogenous levels of total MVP.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	MVP

## **Target Details**

Alternative Name: MVP (MVP Products)  Background: Description: Required for normal vault structure. Vaults are multi-subunit structures that may act as scaffolds for proteins involved in signal transduction. Vaults may also play a role in nucleo-cytoplasmic transport. Down-regulates IFNG-mediated STAT1 signaling and subsequence activation of JAK. Down-regulates SRC activity and signaling through MAP kinases.  Gene: MVP  Molecular Weight: 99kDa		
act as scaffolds for proteins involved in signal transduction. Vaults may also play a role in nucleo-cytoplasmic transport. Down-regulates IFNG-mediated STAT1 signaling and subseq activation of JAK. Down-regulates SRC activity and signaling through MAP kinases.  Gene: MVP	Alternative Name:	cts)
nucleo-cytoplasmic transport. Down-regulates IFNG-mediated STAT1 signaling and subseq activation of JAK. Down-regulates SRC activity and signaling through MAP kinases.  Gene: MVP	Background:	
activation of JAK. Down-regulates SRC activity and signaling through MAP kinases.  Gene: MVP		
Gene: MVP		iic transport. Down-regulates IFNG-mediated STAT1 signaling and subsequent
		Down-regulates SRC activity and signaling through MAP kinases.
Molecular Weight: 99kDa		
	Molecular Weight:	
Gene ID: 9961	Sene ID:	
UniProt: Q14764	JniProt:	
Application Details	Application Details	

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months


#### **Publications**

Product cited in:

Qiu, Li, Kong, Li, Niu, Pan: "Herbal SGR Formula Prevents Acute Ethanol-Induced Liver Steatosis via Inhibition of Lipogenesis and Enhancement Fatty Acid Oxidation in Mice." in: **Evidence-based complementary and alternative medicine : eCAM**, Vol. 2015, pp. 613584, (2015) (


#### PubMed).

#### **Images**



#### **Immunohistochemistry**

**Image 1.** ABIN6276995 at 1/100 staining Rat brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at  $22_{i}$ ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary



#### **Western Blotting**

**Image 2.** Western blot analysis of Hela whole cell lysates, using MVP Antibody. The lane on the left is treated with the antigen-specific peptide.