



[Go to Product page](#)

Datasheet for ABIN6263459  
**anti-MYCBP antibody (N-Term)**

2 Images

Overview

Quantity:	100 µL
Target:	MYCBP
Binding Specificity:	N-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MYCBP antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human MYCBP, corresponding to a region within N-terminal amino acids.
Isotype:	IgG
Specificity:	MYCBP Antibody detects endogenous levels of total MYCBP.
Predicted Reactivity:	Bovine,Rabbit,Dog,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	MYCBP
---------	-------

## Target Details

---

Alternative Name:	MYCBP ( <a href="#">MYCBP Products</a> )
Background:	Description: May control the transcriptional activity of MYC. Stimulates the activation of E box-dependent transcription by MYC. Gene: MYCBP
Molecular Weight:	12 kDa
Gene ID:	26292
UniProt:	<a href="#">Q99417</a>

## Application Details


---

Application Notes:	WB 1:1000-3000, IHC 1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

## Handling


---

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



### Immunohistochemistry

**Image 1.** ABIN6272552 at 1/100 staining Mouse pancreas tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



### Western Blotting

**Image 2.** Western blot analysis of MYCBP using HeLa whole cell lysates