



[Go to Product page](#)

Datasheet for ABIN6263498
anti-MYCN antibody (Internal Region)

2 Images

Overview

Quantity:	100 µL
Target:	MYCN
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MYCN antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human n-Myc, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	N-Myc Antibody detects endogenous levels of total n-Myc.
Predicted Reactivity:	Pig,Bovine,Sheep,Chicken
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	MYCN
---------	------

Target Details


Alternative Name:	MYCN (MYCN Products)
Background:	Description: Positively regulates the transcription of MYCNOS in neuroblastoma cells. Gene: MYCN
Molecular Weight:	49 kDa
Gene ID:	4613
UniProt:	P04198

Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



Western Blotting

Image 1. Western blot analysis of N Myc expression in CACO-2 cells



Immunohistochemistry

Image 2. ABIN6268752 at 1/100 staining human brain tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.