.-online.com antibodies

Datasheet for ABIN6263876 anti-OR52K2 antibody

2 Images



Overview

Quantity:	100 µL
Target:	OR52K2
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OR52K2 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC), Western Blotting (WB)

Product Details


Immunogen:	A synthesized peptide derived from human OR52K2
Isotype:	IgG
Specificity:	OR52K2 Antibody detects endogenous levels of total OR52K2
Cross-Reactivity:	Human
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	OR52K2
Alternative Name:	OR52K2 (OR52K2 Products)
Background:	Description: Odorant receptor.


Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6263876 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
	Gene: OR52K2
Molecular Weight:	35 kDa
Gene ID:	119774
UniProt:	Q8NGK3
Application Details	
Application Notes:	WB 1:1000-3000 IHC 1:200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , $$ pH 7.4, 150 $$ mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C.Stable for 12 months from date of receipt
Expiry Date:	12 months



Western Blotting

Image 1. Western blot analysis OR52K2 using HT-29 whole cell lysates



Immunohistochemistry

Image 2. ABIN6272759 at 1/100 staining Human liver cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.