# antibodies - online.com







# anti-PAFAH2 antibody (Internal Region)

**Images** 



$\sim$	
( )\/\Di	view
	VICVV

Quantity:	100 μL
Target:	PAFAH2
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PAFAH2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

#### **Product Details**

Immunogen:	A synthesized peptide derived from human PAFAH2, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	PAFAH2 Antibody detects endogenous levels of total PAFAH2.
Predicted Reactivity:	Pig,Horse,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

#### **Target Details**

Target:	PAFAH2	

## **Target Details**


Alternative Name:	PAFAH2 (PAFAH2 Products)
Background:	Description: Has a marked selectivity for phospholipids with short acyl chains at the sn-2 position. May share a common physiologic function with the plasma-type enzyme.  Gene: PAFAH2
Molecular Weight:	44-46 kDa
Gene ID:	5051
UniProt:	Q99487

# Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500
Restrictions:	For Research Use only


# Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit lgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



## Immunohistochemistry

**Image 1.** ABIN6273286 at 1/100 staining Human pancreas cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



### Western Blotting

**Image 2.** Western blot analysis of extracts of mouse liver tissue, using PAFAH2 antibody.