# antibodies - online.com







## anti-PATE3 antibody (Internal Region)

**Images** 

Overview

Target Details

PATE3

Target:



Quantity:	100 μL
Target:	PATE3
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PATE3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA
Product Details	
Immunogen:	A synthesized peptide derived from human PATE3, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	PATE3 Antibody detects endogenous levels of total PATE3.
Predicted Reactivity:	Pig,Sheep,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling

#### **Target Details**


Alternative Name:	PATE3 (PATE3 Products)
Background:	Gene: PATE3
Molecular Weight:	12kDa
Gene ID:	100169851
UniProt:	B3GLJ2

## **Application Details**

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



#### **Immunohistochemistry**

**Image 1.** ABIN6277186 at 1/100 staining Mouse testis tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at  $22_{\rm i}$  aC. An HRP conjugated goat anti-rabbit antibody was used as the secondary



#### **Western Blotting**

**Image 2.** Western blot analysis of extracts of HeLa, using PATE3 antibody. The lane on the left is treated with the antigen-specific peptide.