antibodies.com

Datasheet for ABIN6264148 anti-PEX12 antibody (Internal Region)

2 Images



Overview

Quantity:	100 µL	
Target:	PEX12	
Binding Specificity:	Internal Region	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PEX12 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF), Immunohistochemistry (IHC)	

Product Details

Immunogen:	A synthesized peptide derived from human PEX12, corresponding to a region within the internal amino acids.
lsotype:	lgG
Specificity:	PEX12 Antibody detects endogenous levels of total PEX12.
Predicted Reactivity:	Horse,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

Т	ar	a	et:

PEX12

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6264148 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details


Alternative Name:	PEX12 (PEX12 Products)
Background:	Description: Required for protein import into peroxisomes. Gene: PEX12
Molecular Weight:	41 kDa
Gene ID:	5193
UniProt:	000623

Application Details

Application Notes:	WB 1:1000-3000, IHC 1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



Immunohistochemistry

Image 1. ABIN6279651 at 1/100 staining Mouse kidney tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22_{j} aC. An HRP conjugated goat anti-rabbit antibody was used as the secondary

kDa 250-150-100-75-50-37-25-20-15-

Western Blotting

Image 2. Western blot analysis of PEX12 using LOVO whole cell lysates

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN6264148 | 09/10/2023 | Copyright antibodies-online. All rights reserved.