

# Datasheet for ABIN6264460

# anti-DLG4 antibody (C-Term)

2 Images



Go to Product page

Overview	
Quantity:	100 μL
Target:	DLG4
Binding Specificity:	C-Term
Reactivity:	Rat, Mouse, Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DLG4 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA
Product Details	
Immunogen:	A synthesized peptide derived from human PSD95, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	PSD95 Antibody detects endogenous levels of total PSD95.
Predicted Reactivity:	Zebrafish,Bovine,Horse,Sheep,Rabbit
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	DLG4


## **Target Details**


DLG4 (DLG4 Products)
Description: Interacts with the cytoplasmic tail of NMDA receptor subunits and shaker-type
potassium channels. Required for synaptic plasticity associated with NMDA receptor signaling
Overexpression or depletion of DLG4 changes the ratio of excitatory to inhibitory synapses in
hippocampal neurons. May reduce the amplitude of ASIC3 acid-evoked currents by retaining
the channel intracellularly. May regulate the intracellular trafficking of ADR1B. Also regulates
AMPA-type glutamate receptor (AMPAR) immobilization at postsynaptic density keeping the
channels in an activated state in the presence of glutamate and preventing synaptic
depression.
Gene: DLG4
105 kDa
1742
P78352
Regulation of Muscle Cell Differentiation, Synaptic Membrane, Skeletal Muscle Fiber
Development, Asymmetric Protein Localization, Regulation of long-term Neuronal Synaptic
Plasticity
WB 1:500-1:2000, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
For Research Use only
Liquid
1 mg/mL
1 mg/mL  Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %
Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %
Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.  Sodium azide
Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.  Sodium azide  This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Expiry Date:

12 months

### **Images**





#### **Immunohistochemistry**

**Image 1.** ABIN6268831 at 1/200 staining human brain tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

#### **Western Blotting**

**Image 2.** Western blot analysis of PSD95 expression in Brain (Rat) Tissue Lysate, The lane on the left is treated with the antigen-specific peptide.