antibodies -online.com





anti-PSMD2 antibody (C-Term)





Go to Product page

| \sim | | | | | | |
|--------|-----|--------|----|-----|-------------|---|
| | 1// | \Box | r۱ | / [| \triangle | ٨ |


| Quantity: | 100 μL |
|----------------------------|--|
| Target: | PSMD2 |
| Binding Specificity: | C-Term |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PSMD2 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |
| | |
| Product Details | |
| Product Details Immunogen: | A synthesized peptide derived from human PSMD2, corresponding to a region within C-terminal amino acids. |
| | |
| Immunogen: | amino acids. |
| Immunogen: Isotype: | amino acids. IgG |

Target Details

Target: PSMD2


Target Details

| Alternative Name: | PSMD2 (PSMD2 Products) | |
|---------------------|---|--|
| Background: | Description: Component of the 26S proteasome, a multiprotein complex involved in the ATP-dependent degradation of ubiquitinated proteins. This complex plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins, which could impair cellular functions, and by removing proteins whose functions are no longer required. Therefore, the proteasome participates in numerous cellular processes, including cell cycle progression, apoptosis, or DNA damage repair. Gene: PSMD2 | |
| Molecular Weight: | 100kDa | |
| Gene ID: | 5708 | |
| UniProt: | Q13200 | |
| Pathways: | Mitotic G1-G1/S Phases, DNA Replication, Synthesis of DNA, Ubiquitin Proteasome Pathway | |
| Application Details | | |
| Application Notes: | WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 mg/mL | |
| Buffer: | Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol. | |
| Preservative: | Sodium azide | |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20 °C. Stable for 12 months from date of receipt. | |
| Expiry Date: | 12 months | |




Western Blotting

Image 1. Western blot analysis of Hela whole cell lysates, using PSMD2 Antibody. The lane on the left is treated with the antigen-specific peptide.



Immunohistochemistry

Image 2. ABIN6277006 at 1/100 staining Human brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22¡ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary



Immunofluorescence (fixed cells)

Image 3. ABIN6277006 staining COLO205 cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25;ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37;ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibod

Please check the product details page for more images. Overall 4 images are available for ABIN6264489.