# antibodies - online.com







## anti-RAD9A antibody (C-Term)





Goto	Product	nage
	iioaact	page

Overview

0.10.11011	
Quantity:	100 μL
Target:	RAD9A
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RAD9A antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)
Product Details	
Immunogen:	A synthesized peptide derived from human RAD9, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	RAD9 Antibody detects endogenous levels of total RAD9.
Predicted Reactivity:	Pig,Zebrafish,Horse,Sheep,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	RAD9A

### **Target Details**


Alternative Name:	RAD9A (RAD9A Products)	
Background:	Description: Component of the 9-1-1 cell-cycle checkpoint response complex that plays a majo role in DNA repair. The 9-1-1 complex is recruited to DNA lesion upon damage by the RAD17-replication factor C (RFC) clamp loader complex. Acts then as a sliding clamp platform on DNA	
	for several proteins involved in long-patch base excision repair (LP-BER). The 9-1-1 complex	
	stimulates DNA polymerase beta (POLB) activity by increasing its affinity for the 3'-OH end of	
	the primer-template and stabilizes POLB to those sites where LP-BER proceeds, endonuclease	
	FEN1 cleavage activity on substrates with double, nick, or gap flaps of distinct sequences and	
	lengths, and DNA ligase I (LIG1) on long-patch base excision repair substrates. The 9-1-1	
	complex is necessary for the recruitment of RHN01 to sites of double-stranded breaks (DSB)	
	occurring during the S phase. RAD9A possesses 3'->5' double stranded DNA exonuclease activity. Its phosphorylation by PRKCD may be required for the formation of the 9-1-1 complex. Gene: RAD9A	
Molecular Weight:	42kDa	
Gene ID:	5883	
UniProt:	Q99638	
Pathways:	Positive Regulation of Response to DNA Damage Stimulus	
Application Details		
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Storage:	-20 °C	

#### Handling

Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.	
------------------	---	--


Expiry Date: 12 months

#### **Images**



#### **Immunohistochemistry**

**Image 1.** ABIN6276941 at 1/100 staining Rat liver tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22<sub>i</sub>aC. An HRP conjugated goat anti-rabbit antibody was used as the secondary



#### **Western Blotting**

**Image 2.** Western blot analysis of extracts of HeLa, using RAD9A antibody. The lane on the left is treated with the antigen-specific peptide.