antibodies -online.com





anti-RIOK3 antibody (Internal Region)

2 Images



Go to Product page

\sim	
()\/\	rview
\cup	

Quantity:	100 μL
Target:	RIOK3
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RIOK3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human RIOK3, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	RIOK3 Antibody detects endogenous levels of total RIOK3.
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Sheep,Rabbit,Dog,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	RIOK3	

Target Details		
Alternative Name:	RIOK3 (RIOK3 Products)	
Background:	Description: Involved in regulation of type I interferon (IFN)-dependent immune response which	
	plays a critical role in the innate immune response against DNA and RNA viruses. May act as an	
	adapter protein essential for the recruitment of TBK1 to IRF3 (PubMed:24807708).	
	Phosphorylates IFIH1 on 'Ser-828' interfering with IFIH1 filament assembly on long dsRNA and	
	resulting in attenuated IFIH1-signaling (PubMed:25865883). Can inhibit CASP10 isoform 7-	
	mediated activation of the NF-kappaB signaling pathway (PubMed:19557502). May play a role	
	in the biogenesis of the 40S ribosomal subunit. Involved in the processing of 21S pre-rRNA to	
	the mature 18S rRNA (PubMed:22418843).	
	Gene: RIOK3	
Molecular Weight:	59 kDa	
Gene ID:	8780	
UniProt:	014730	
Application Details		
Application Notes:	WB 1:1000-3000, IF/ICC 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %	
	glycerol.	
Preservative:	Sodium azide	

Store at -20 °C. Stable for 12 months from date of receipt.

should be handled by trained staff only.

-20 °C

12 months


This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which


Precaution of Use:

Storage Comment:

Storage:

Expiry Date:





Immunofluorescence (fixed cells)

Image 1. ABIN6278629 staining HepG2 cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibod

Western Blotting

Image 2. Western blot analysis of RIOK3 using K562 whole cell lysates