antibodies -online.com





anti-RNF41 antibody (Internal Region)

3 Im





	Go to Product page

()	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	$I \vee I$	ew

Quantity:	100 μL
Target:	RNF41
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RNF41 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunocytochemistry (ICC),
	Immunofluorescence (IF)
Product Details	
Immunogen:	A synthesized peptide derived from human RNF41, corresponding to a region within the internal
	amino acids.
Isotype:	IgG
Specificity:	RNF41 Antibody detects endogenous levels of total RNF41.
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Sheep,Rabbit,Dog,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling
	Resin (Thermo Fisher Scientific).

Target Details

Target: RNF41


Target Details

Alternative Name:	RNF41 (RNF41 Products)
Background:	Description: Acts as E3 ubiquitin-protein ligase and regulates the degradation of target proteins
	Polyubiquitinates MYD88. Negatively regulates MYD88-dependent production of
	proinflammatory cytokines. Can promote TRIF-dependent production of type I interferon and
	inhibits infection with vesicular stomatitis virus (By similarity). Promotes also activation of
	TBK1 and IRF3. Involved in the ubiquitination of erythropoietin (EPO) and interleukin-3 (IL-3)
	receptors. Thus, through maintaining basal levels of cytokine receptors, RNF41 is involved in
	the control of hematopoietic progenitor cell differentiation into myeloerythroid lineages (By
	similarity). Contributes to the maintenance of steady-state ERBB3 levels by mediating its
	growth factor-independent degradation. Involved in the degradation of the inhibitor of apoptosi
	BIRC6 and thus is an important regulator of cell death by promoting apoptosis. Acts also as a
	PRKN modifier that accelerates its degradation, resulting in a reduction of PRKN activity,
	influencing the balance of intracellular redox state. The RNF41-PRKN pathway regulates
	autophagosome-lysosome fusion during late mitophagy. Mitophagy is a selective form of
	autophagy necessary for mitochondrial quality control (PubMed:24949970).
	Gene: RNF41
Molecular Weight:	36 kDa
Gene ID:	10193
UniProt:	Q9H4P4
Pathways:	SARS-CoV-2 Protein Interactome
Application Details	
Application Notes:	WB 1:1000-3000, IHC 1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %
	glycerol.
Preservative:	Sodium azide

Handling


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.	
Expiry Date:	12 months	

Images




Western Blotting

Image 1. Western blot analysis of RNF41 using HeLa whole cell lysates



Immunohistochemistry

Image 2. ABIN6279398 at 1/100 staining Human prostate tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at $22_{\rm i}$ aC. An HRP conjugated goat anti-rabbit antibody was used as the secondary



Western Blotting

Image 3. Western blot analysis of extracts from rat muscle, using RNF41 Antibody.