

Datasheet for ABIN6264963  
**anti-SEC63 antibody (C-Term)**[Go to Product page](#)

## 3 Images

## Overview

Quantity:	100 µL
Target:	SEC63
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SEC63 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

## Product Details

Immunogen:	A synthesized peptide derived from human SEC63, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	SEC63 Antibody detects endogenous levels of total SEC63.
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Sheep,Rabbit,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

## Target Details

Target:	SEC63
---------	-------

## Target Details


Alternative Name:	SEC63 ( <a href="#">SEC63 Products</a> )
Background:	Description: Required for integral membrane and secreted preprotein translocation across the endoplasmic reticulum membrane. Gene: SEC63
Molecular Weight:	83 kDa
Gene ID:	11231
UniProt:	<a href="#">Q9UGP8</a>
Pathways:	<a href="#">ER-Nucleus Signaling</a>

## Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months




Immunohistochemistry

**Image 1.** ABIN6273315 at 1/100 staining Human thyroid cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



Western Blotting

**Image 2.** Western blot analysis of extracts of PC-3 cells, using SEC63 antibody.



Western Blotting

**Image 3.** Western blot analysis of extracts from 3t3, using SEC63 Antibody. The lane on the left was treated with blocking peptide.