## antibodies -online.com





## anti-SETD7 antibody (Internal Region)



Image



Go to Product page


Overview	
Quantity:	100 μL
Target:	SETD7
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SETD7 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)
Product Details	
Immunogen:	A synthesized peptide derived from human SETD7, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	SETD7 Antibody detects endogenous levels of total SETD7.
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Sheep,Rabbit,Dog,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	SETD7

## **Target Details**

Expiry Date:

· g	
Alternative Name:	SETD7 (SETD7 Products)
Background:	Description: Histone methyltransferase that specifically monomethylates 'Lys-4' of histone H3.
	H3 'Lys-4' methylation represents a specific tag for epigenetic transcriptional activation. Plays a
	central role in the transcriptional activation of genes such as collagenase or insulin. Recruited
	by IPF1/PDX-1 to the insulin promoter, leading to activate transcription. Has also
	methyltransferase activity toward non-histone proteins such as p53/TP53, TAF10, and possibly
	TAF7 by recognizing and binding the [KR]-[STA]-K in substrate proteins. Monomethylates 'Lys-
	189' of TAF10, leading to increase the affinity of TAF10 for RNA polymerase II. Monomethylates
	'Lys-372' of p53/TP53, stabilizing p53/TP53 and increasing p53/TP53-mediated transcriptional
	activation.
	Gene: SETD7
Molecular Weight:	41 kDa
Gene ID:	80854
UniProt:	Q8WTS6
A 1: 1: D 1:1	
Application Details	
Application Notes:	WB 1:1000-3000, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %
	glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.

12 months



## Western Blotting

Image 1. Western blot analysis SETD7 using Jurkat whole cell lysates