antibodies -online.com







anti-PTPN11 antibody (C-Term)





\sim	
()\/\	rview
\circ	

Quantity:	100 μL
Target:	PTPN11
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PTPN11 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human SHP2, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	SHP2 Antibody detects endogenous levels of total SHP2.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog,Chicken
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	PTPN11	

Target Details


rarget betano				
Alternative Name:	PTPN11 (PTPN11 Products)			
Target Type:	Viral Protein			
Background:	Description: Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus. Positively regulates MAPK signal transduction pathway (PubMed:28074573). Dephosphorylates GAB1, ARHGAP35 and EGFR (PubMed:28074573). Dephosphorylates ROCK2 at 'Tyr-722' resulting in stimulatation of its RhoA binding activity. Dephosphorylates CDC73 (PubMed:26742426). Gene: PTPN11			
Molecular Weight:	70kDa			
Gene ID:	5781			
UniProt:	Q06124			
Pathways:	JAK-STAT Signaling, RTK Signaling, TCR Signaling, Interferon-gamma Pathway, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Negative Regulation of Hormone Secretion, Carbohydrate Homeostasis, Toll-Like Receptor Cascades, CXCR4-mediated Signaling Events, Signaling Events mediated by VEGFR1 and VEGFR2, Signaling of Hepatocyte Growth Factor Receptor, VEGFR1 Specific Signals, BCR Signaling, Warburg Effect			
Application Details				
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000			
Restrictions:	For Research Use only			
Handling				
Format:	Liquid			
Concentration:	1 mg/mL			
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.			
Preservative:	Sodium azide			
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.			
Storage:	-20 °C			

Handling

Storage Comment:	Store at -20 °C	. Stable for 12	2 months from	date of receipt.


Expiry Date: 12 months

Images




Immunofluorescence (fixed cells)

Image 1. ABIN6269332 staining A431 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



Western Blotting

Image 2. Western blot analysis of SHP-2 expression in A431 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



Immunohistochemistry

Image 3. ABIN6269332 at 1/100 staining Mouse muscle tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

Please check the product details page for more images. Overall 4 images are available for ABIN6265075.