antibodies .- online.com







anti-SMARCE1 antibody (C-Term)

Images



\sim				
	$ V \cap$	r\/I	19	٨

Quantity:	100 μL	
Target:	SMARCE1	
Binding Specificity:	C-Term	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SMARCE1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)	
Product Details		

Immunogen:	A synthesized peptide derived from human SMARCE1, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	SMARCE1 Antibody detects endogenous levels of total SMARCE1.
Predicted Reactivity:	Pig,Horse,Sheep,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	SMARCE1	


Target Details

Alternative Name:	SMARCE1 (SMARCE1 Products)
Background:	Description: Involved in transcriptional activation and repression of select genes by chromatin
	remodeling (alteration of DNA-nucleosome topology). Component of SWI/SNF chromatin
	remodeling complexes that carry out key enzymatic activities, changing chromatin structure by
	altering DNA-histone contacts within a nucleosome in an ATP-dependent manner. Belongs to
	the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the
	neuron-specific chromatin remodeling complex (nBAF complex). During neural development a
	switch from a stem/progenitor to a postmitotic chromatin remodeling mechanism occurs as
	neurons exit the cell cycle and become committed to their adult state. The transition from
	proliferating neural stem/progenitor cells to postmitotic neurons requires a switch in subunit
	composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and
	differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and
	PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and
	DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF
	complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem
	cells. The nBAF complex along with CREST plays a role regulating the activity of genes
	essential for dendrite growth (By similarity). Required for the coactivation of estrogen
	responsive promoters by SWI/SNF complexes and the SRC/p160 family of histone
	acetyltransferases (HATs). Also specifically interacts with the CoREST corepressor resulting in
	repression of neuronal specific gene promoters in non-neuronal cells.
	Gene: SMARCE1
Molecular Weight:	46kDa
Gene ID:	6605
UniProt:	Q969G3
Pathways:	Chromatin Binding
Application Details	
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
	Liquid

Handling


Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. ABIN6277544 at 1/100 staining Mouse liver tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22¡ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary



Western Blotting

Image 2. Western blot analysis of Hela whole cell lysates, using SMARCE1 Antibody. The lane on the left is treated with the antigen-specific peptide.