

Datasheet for ABIN6265404
anti-Syntaxin 16 antibody (N-Term)

2 Images

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Syntaxin 16 (STX16)
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Syntaxin 16 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Immunogen:	A synthesized peptide derived from human Syntaxin 16, corresponding to a region within N-terminal amino acids.
Isotype:	IgG
Specificity:	Syntaxin 16 Antibody detects endogenous levels of total Syntaxin 16.
Predicted Reactivity:	Bovine,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	Syntaxin 16 (STX16)
---------	---------------------

Target Details


Alternative Name:	STX16 (STX16 Products)
Background:	Description: SNARE involved in vesicular transport from the late endosomes to the trans-Golgi network. Gene: STX16
Molecular Weight:	42 kDa
Gene ID:	8675
UniProt:	014662
Pathways:	Synaptic Vesicle Exocytosis

Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



Western Blotting

Image 1. Western blot analysis of extracts from Jurkat, using Syntaxin 16 antibody.



Immunohistochemistry

Image 2. ABIN6273174 at 1/100 staining Human lymph cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.