antibodies - online.com







anti-TAL1 antibody (C-Term)





Overview	
Quantity:	100 μL
Target:	TAL1
Binding Specificity:	C-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TAL1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
lmmunogen:	A synthesized peptide derived from human Tal1, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	Tal1 Antibody detects endogenous levels of total Tal1.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	TAL1

Target Details


Alternative Name:	TAL1 (TAL1 Products)
Background:	Description: Implicated in the genesis of hemopoietic malignancies. It may play an important
	role in hemopoietic differentiation. Serves as a positive regulator of erythroid differentiation (By
	similarity).
	Gene: TAL1
Molecular Weight:	34 kDa
Gene ID:	6886
UniProt:	P17542
Pathways:	Stem Cell Maintenance

Application Details

Application Notes:	WB 1:1000-3000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



Western Blotting

Image 1. Western blot analysis of TAL-1 using Jurkat whole cell lysates



Immunofluorescence (fixed cells)

Image 2. ABIN6278639 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod