

Datasheet for ABIN6265487
anti-T-Box 15/18 antibody

3 Images

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	T-Box 15/18 (TBX15/18)
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This T-Box 15/18 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human TBX15/18
Isotype:	IgG
Specificity:	TBX15/18 Antibody detects endogenous levels of total TBX15/18
Cross-Reactivity:	Human, Mouse (Murine), Rat (Rattus)
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	T-Box 15/18 (TBX15/18)
Alternative Name:	TBX15/18 (TBX15/18 Products)

Target Details


Background:	Description: Acts as transcriptional repressor involved in developmental processes of a variety of tissues and organs, including the heart and coronary vessels, the ureter and the vertebral column. Required for embryonic development of the sino atrial node (SAN) head area. Gene: TBX18
Molecular Weight:	65 kDa
Gene ID:	9096
UniProt:	O95935 , Q96SF7

Application Details

Application Notes:	WB 1:1000-3000 IHC 1:200, IF/ICC 1:100-1:500
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C.Stable for 12 months from date of receipt
Expiry Date:	12 months




Immunohistochemistry

Image 1. ABIN6278641 at 1/100 staining Rat brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary



Western Blotting

Image 2. Western blot analysis of TBX15/18 using HepG2 whole cell lysates



Immunofluorescence (fixed cells)

Image 3. ABIN6278641 staining HepG2 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody