# antibodies - online.com







## anti-Tropomodulin 1 antibody (N-Term)

**Images** 



$\sim$	
( )\/户	rview
$\circ$	V I C V V

Target:

100 μL
Tropomodulin 1 (TMOD1)
N-Term
Human, Mouse, Rat
Rabbit
Polyclonal
This Tropomodulin 1 antibody is un-conjugated
Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF),
Immunocytochemistry (ICC)
A synthesized peptide derived from human TMOD1, corresponding to a region within N-terminal
amino acids.
IgG
TMOD1 Antibody detects endogenous levels of total TMOD1.
Bovine,Horse,Sheep,Rabbit,Dog,Chicken
The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling
Resin (Thermo Fisher Scientific).


Tropomodulin 1 (TMOD1)

### **Target Details**

Expiry Date:


12 months

rarget Details	
Alternative Name:	TMOD1 (TMOD1 Products)
Background:	Description: Blocks the elongation and depolymerization of the actin filaments at the pointed end. The Tmod/TM complex contributes to the formation of the short actin protofilament, which in turn defines the geometry of the membrane skeleton. May play an important role in regulating the organization of actin filaments by preferentially binding to a specific tropomyosin isoform at its N-terminus.  Gene: TMOD1
Molecular Weight:	41 kDa
Gene ID:	7111
UniProt:	P28289
Application Details	
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.




#### **Immunohistochemistry**

**Image 1.** ABIN6273177 at 1/100 staining Human lymph cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



#### **Western Blotting**

**Image 2.** Western blot analysis of extracts of human brain tissue, using TMOD1 antibody.



#### **Western Blotting**

**Image 3.** Western blot analysis of extracts from Mouse brain, using TMOD1 Antibody. The lane on the left was treated with blocking peptide.