antibodies -online.com





anti-TNFAIP3 antibody (Internal Region)





Go to Product page

_				
(11/0	K\ /	$ \Delta $	Α.
	\cup	1 \/	ı⊢۱	Λ

Target:

Quantity:	100 μL
Target:	TNFAIP3
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TNFAIP3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized peptide derived from human A20/TNFAIP3, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	A20/TNFAIP3 Antibody detects endogenous levels of total A20/TNFAIP3.
Predicted Reactivity:	Pig,Bovine,Horse,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).
Target Details	

TNFAIP3

Target Details	
Alternative Name:	TNFAIP3 (TNFAIP3 Products)
Background:	Description: Ubiquitin-editing enzyme that contains both ubiquitin ligase and deubiquitinase
	activities. Involved in immune and inflammatory responses signaled by cytokines, such as TNF-
	alpha and IL-1 beta, or pathogens via Toll-like receptors (TLRs) through terminating NF-kappa-B
	activity. Essential component of a ubiquitin-editing protein complex, comprising also RNF11,
	ITCH and TAX1BP1, that ensures the transient nature of inflammatory signaling pathways. In
	cooperation with TAX1BP1 promotes disassembly of E2-E3 ubiquitin protein ligase complexes
	in IL-1R and TNFR-1 pathways, affected are at least E3 ligases TRAF6, TRAF2 and BIRC2, and
	E2 ubiquitin-conjugating enzymes UBE2N and UBE2D3. In cooperation with TAX1BP1 promotes
	ubiquitination of UBE2N and proteasomal degradation of UBE2N and UBE2D3. Upon TNF
	stimulation, deubiquitinates 'Lys-63'-polyubiquitin chains on RIPK1 and catalyzes the formation
	of 'Lys-48'-polyubiquitin chains. This leads to RIPK1 proteasomal degradation and consequently
	termination of the TNF- or LPS-mediated activation of NF-kappa-B. Deubiquitinates TRAF6
	probably acting on 'Lys-63'-linked polyubiquitin. Upon T-cell receptor (TCR)-mediated T-cell
	activation, deubiquitinates 'Lys-63'-polyubiquitin chains on MALT1 thereby mediating
	disassociation of the CBM (CARD11:BCL10:MALT1) and IKK complexes and preventing
	sustained IKK activation. Deubiquitinates NEMO/IKBKG, the function is facilitated by TNIP1 and
	leads to inhibition of NF-kappa-B activation. Upon stimulation by bacterial peptidoglycans,
	probably deubiquitinates RIPK2. Can also inhibit I-kappa-B-kinase (IKK) through a non-catalytic
	mechanism which involves polyubiquitin, polyubiquitin promotes association with IKBKG and
	prevents IKK MAP3K7-mediated phosphorylation. Targets TRAF2 for lysosomal degradation. In
	vitro able to deubiquitinate 'Lys-11'-, 'Lys-48'- and 'Lys-63' polyubiquitin chains. Inhibitor of
	programmed cell death. Has a role in the function of the lymphoid system. Required for LPS-
	induced production of proinflammatory cytokines and IFN beta in LPS-tolerized macrophages.
	Gene: TNFAIP3
Molecular Weight:	90kDa
Gene ID:	7128
UniProt:	P21580
Pathways:	TLR Signaling, Activation of Innate immune Response, Cellular Response to Molecule of
	Bacterial Origin, Production of Molecular Mediator of Immune Response

Application Details

Application Notes: WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000


Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling


Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

Images




Immunohistochemistry

Image 1. ABIN6277109 at 1/100 staining Rat kidney tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at $22_{\rm i}$ aC. An HRP conjugated goat anti-rabbit antibody was used as the secondary



Western Blotting

Image 2. Western blot analysis of Hepg2 whole cell lysates, using TNFAIP3 Antibody. The lane on the left is treated with the antigen-specific peptide.



Western Blotting

Image 3. Western blot analysis of extracts from various samples, using TNFAIP3 Antibody. Lane 1: Hela treated with blocking peptide; Lane 2: Hela; Lane 3: HepG2.

Please check the product details page for more images. Overall 4 images are available for ABIN6265632.