# antibodies -online.com





# anti-TRADD antibody

3 Images



Go to Product page

# Overview

Quantity:	100 μL
Target:	TRADD
Reactivity:	Human, Mouse, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRADD antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

# **Product Details**

Immunogen:	A synthesized peptide derived from human TRADD
Isotype:	IgG
Specificity:	TRADD antibody detects endogenous levels of total TRADD
Cross-Reactivity:	Human, Monkey, Mouse (Murine)
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).


# **Target Details**

Target:	TRADD
Alternative Name:	TRADD (TRADD Products)

# **Target Details**


rarget Details	
Background:	Description: The nuclear form acts as a tumor suppressor by preventing ubiquitination and
	degradation of isoform p19ARF/ARF of CDKN2A by TRIP12: acts by interacting with TRIP12,
	leading to disrupt interaction between TRIP12 and isoform p19ARF/ARF of CDKN2A (By
	similarity). Adapter molecule for TNFRSF1A/TNFR1 that specifically associates with the
	cytoplasmic domain of activated TNFRSF1A/TNFR1 mediating its interaction with FADD.
	Overexpression of TRADD leads to two major TNF-induced responses, apoptosis and activation
	of NF-kappa-B.
	Gene: TRADD
Molecular Weight:	34kDa
Gene ID:	8717
UniProt:	Q15628
Pathways:	NF-kappaB Signaling, Apoptosis, Caspase Cascade in Apoptosis, Positive Regulation of
	Endopeptidase Activity, Hepatitis C
Application Details	
Application Notes:	WB: 1:500~1:3000 IHC: 1:50~1:200, IF/ICC 1:100-1:500
Restrictions:	For Research Use only
Handling	

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C.Stable for 12 months from date of receipt
Expiry Date:	12 months




### Immunofluorescence (fixed cells)

**Image 1.** ABIN6266665 staining COS7 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



## **Immunohistochemistry**

**Image 2.** ABIN6266665 at 1/100 staining human brain tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



# **Western Blotting**

**Image 3.** Western blot analysis on COS7 and HuvEc cell lysate using TRADD Antibody. The lane on the left is treated with the antigen-specific peptide.