

Datasheet for ABIN6265740
anti-TRIM44 antibody (Internal Region)

3 Images

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	TRIM44
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRIM44 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human TRIM44, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	TRIM44 Antibody detects endogenous levels of total TRIM44.
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Sheep,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	TRIM44
---------	--------

Target Details


Alternative Name:	TRIM44 (TRIM44 Products)
Background:	Description: May play a role in the process of differentiation and maturation of neuronal cells (By similarity). May regulate the activity of TRIM17. Is a negative regulator of PAX6 expression (PubMed:26394807). Gene: TRIM44
Molecular Weight:	50-55kDa
Gene ID:	54765
UniProt:	Q96DX7

Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months




Western Blotting

Image 1. Western blot analysis of extracts of human liver tissue, using TRIM44 antibody.



Immunohistochemistry

Image 2. ABIN6273021 at 1/100 staining Human liver tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



Western Blotting

Image 3. Western blot analysis of extracts from various samples, using TRIM44 antibody. Lane 1: rat heart treated with blocking peptide. Lane 2: Rat heart; Lane 3: Hepg2;