# antibodies -online.com





### anti-TYROBP antibody (C-Term)

2 Images

Overview

Binding Specificity:



Go to Product page

# Quantity: 100 μL Target: TYROBP

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Conjugate: This TYROBP antibody is un-conjugated

C-Term

Application: Western Blotting (WB), Immunohistochemistry (IHC), ELISA

#### **Product Details**

Immunogen:	A synthesized peptide derived from human TYROBP, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	TYROBP Antibody detects endogenous levels of total TYROBP.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

#### **Target Details**

Target: TYROBP

#### **Target Details**


Alternative Name:	TYROBP (TYROBP Products)
Background:	Description: Non-covalently associates with activating receptors of the CD300 family. Cross-linking of CD300-TYROBP complexes results in cellular activation. Involved for instance in neutrophil activation mediated by integrin.  Gene: TYROBP
Molecular Weight:	12kDa
Gene ID:	7305
UniProt:	043914

### Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


#### Handling

Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.	
Expiry Date:	12 months	



## Immunohistochemistry

**Image 1.** ABIN6277559 at 1/100 staining Human Head and neck cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22¡ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary



#### **Western Blotting**

**Image 2.** Western blot analysis of extracts of SK-SEP1, using TYROBP antibody. The lane on the left is treated with the antigen-specific peptide.