antibodies -online.com





anti-ZBTB33 antibody (N-Term)






Go to Product page

| \sim | | | | | | | |
|--------|-----|--------|---------------|-----|------------|---|----|
| | 1// | \Box | $r \setminus$ | / [| \bigcirc | 1 | ٨, |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | ZBTB33 |
| Binding Specificity: | N-Term |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ZBTB33 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA |
| Product Details | |
| Immunogen: | A synthesized peptide derived from human ZBTB33, corresponding to a region within N-terminal amino acids. |
| Isotype: | IgG |
| Specificity: | ZBTB33 Antibody detects endogenous levels of total ZBTB33. |
| Purification: | The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific). |
| Target Details | |
| Target: | ZBTB33 |
| Alternative Name: | ZBTB33 (ZBTB33 Products) |

| Target Details | |
|---------------------|--|
| Background: | Description: Transcriptional regulator with bimodal DNA-binding specificity. Binds to methylated |
| | CpG dinucleotides in the consensus sequence 5'-CGCG-3' and also binds to the non-methylated |
| | consensus sequence 5'-CTGCNA-3' also known as the consensus kaiso binding site (KBS). |
| | Recruits the N-CoR repressor complex to promote histone deacetylation and the formation of |
| | repressive chromatin structures in target gene promoters. May contribute to the repression of |
| | target genes of the Wnt signaling pathway. May also activate transcription of a subset of target |
| | genes by the recruitment of CTNND2. Represses expression of MMP7 in conjunction with |
| | transcriptional corepressors CBFA2T3, CBFA2T2 and RUNX1T1 (PubMed:23251453). |
| | Gene: ZBTB33 |
| Molecular Weight: | 74 kDa |
| Gene ID: | 10009 |
| UniProt: | Q86T24 |
| Application Details | |
| Application Notes: | WB 1:1000-3000, ELISA(peptide) 1:20000-1:40000 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 mg/mL |
| Buffer: | Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20 °C. Stable for 12 months from date of receipt. |
| Expiry Date: | 12 months |



Western Blotting

Image 1. Western blot analysis of ZBTB33 using COLO205 whole cell lysates