

Datasheet for ABIN6266119

**anti-ZMYM4 antibody (Internal Region)**[Go to Product page](#)**2** Images

## Overview

Quantity:	100 µL
Target:	ZMYM4
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZMYM4 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA

## Product Details

Immunogen:	A synthesized peptide derived from human ZNF262, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	ZNF262 Antibody detects endogenous levels of total ZNF262.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog,Chicken
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

## Target Details

Target:	ZMYM4
---------	-------

## Target Details


Alternative Name:	ZMYM4 ( <a href="#">ZMYM4 Products</a> )
Background:	Description: Plays a role in the regulation of cell morphology and cytoskeletal organization. Gene: ZMYM4
Molecular Weight:	175kDa
Gene ID:	9202
UniProt:	<a href="#">Q5VZL5</a>

## Application Details

Application Notes:	WB 1:1000, IHC 1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



### Western Blotting

**Image 1.** Western blot analysis of ZNF262 using HepG2 whole cell lysates



### Immunohistochemistry

**Image 2.** ABIN6272467 at 1/100 staining Mouse heart tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.