# antibodies - online.com







# anti-ZNF785 antibody

**Images** 



$\sim$	
( )VPI	view
0 1 0	VICVV

Quantity:	100 μL
Target:	ZNF785
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF785 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF)

#### **Product Details**

Immunogen:	A synthesized peptide derived from human ZNF785
Isotype:	IgG
Specificity:	ZNF785 Antibody detects endogenous levels of total ZNF785
Cross-Reactivity:	Human, Mouse (Murine), Rat (Rattus)
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling
	Resin (Thermo Fisher Scientific).

### **Target Details**

Target:	ZNF785
Alternative Name:	ZNF785 (ZNF785 Products)
Background:	Description: May be involved in transcriptional regulation.

## **Target Details**


	Gene: ZNF785
Molecular Weight:	87kDa
Gene ID:	146540
UniProt:	A8K8V0

# Application Details

Application Notes:	WB 1:1000, IF/ICC 1:100-1:500
Restrictions:	For Research Use only


# Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C.Stable for 12 months from date of receipt
Expiry Date:	12 months



#### **Western Blotting**

**Image 1.** Western blot analysis of ZNF785 using HepG2 whole cell lysates



#### Immunofluorescence (fixed cells)

**Image 2.** ABIN6272469 staining HepG2 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.