

Datasheet for ABIN6266143  
**anti-Slc30a3 antibody (C-Term)**

## 3 Images

[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	Slc30a3
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Slc30a3 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunocytochemistry (ICC), Immunofluorescence (IF)

## Product Details

Immunogen:	A synthesized peptide derived from human ZnT3, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	ZnT3 Antibody detects endogenous levels of total ZnT3.
Predicted Reactivity:	Pig,Bovine,Sheep,Rabbit
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

## Target Details

Target:	Slc30a3
---------	---------

## Target Details


Alternative Name:	SLC30A3 ( <a href="#">Slc30a3 Products</a> )
Background:	Description: Involved in accumulation of zinc in synaptic vesicles. Gene: SLC30A3
Molecular Weight:	42 kDa,48 kDa
Gene ID:	7781
UniProt:	<a href="#">Q99726</a>
Pathways:	<a href="#">Transition Metal Ion Homeostasis</a>

## Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



**Immunohistochemistry**

**Image 1.** ABIN6273188 at 1/100 staining Human lymph cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



**Western Blotting**

**Image 2.** Western blot analysis of extracts of mouse brain tissue, using ZnT3 antibody.

**Western Blotting**

**Image 3.** Western blot analysis of extracts from Mouse brain, using ZnT3 Antibody. The lane on the left was treated with blocking peptide.