# antibodies -online.com







## anti-CD90 antibody (Internal Region)



**Images** 



$\sim$						
	1//	Д	r۱	1	<b>Θ</b> 1	٨

Quantity:	100 μL
Target:	CD90 (THY1)
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CD90 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), ELISA, Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized pentide derived from human THY1 corresponding to a region within the internal

Immunogen:	A synthesized peptide derived from human THY1, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	THY1 Antibody detects endogenous levels of total THY1.
Predicted Reactivity:	Pig,Horse,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

### Target Details

Target:	CD90 (THY1)		

### **Target Details**


Alternative Name:	THY1 (THY1 Products)
Background:	Description: May play a role in cell-cell or cell-ligand interactions during synaptogenesis and other events in the brain.  Gene: THY1
Molecular Weight:	18kDa
Gene ID:	7070
UniProt:	P04216
Pathways:	Cell-Cell Junction Organization

## **Application Details**

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months




#### **Western Blotting**

**Image 1.** Western blot analysis of extracts of fetal brain, using THY1 antibody. The lane on the left is treated with the antigen-specific peptide.



#### Immunofluorescence (fixed cells)

Image 2. ABIN6277108 staining HuvEc cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibod



#### **Immunohistochemistry**

**Image 3.** ABIN6277108 at 1/100 staining Human liver cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22¡ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary