antibodies - online.com







anti-TRIM63 antibody (N-Term)

Images



\sim	
()\/\p	rview
\circ	

Quantity:	100 μL
Target:	TRIM63
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRIM63 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized peptide derived from human TRIM63, corresponding to a region within N-terminal amino acids.
Isotype:	IgG
Specificity:	TRIM63 Antibody detects endogenous levels of total TRIM63.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog

Target Details

Purification:


Target: TRIM63

Resin (Thermo Fisher Scientific).

The antiserum was purified by peptide affinity chromatography using SulfoLinkTM Coupling


Target Details

Alternative Name:	TRIM63 (TRIM63 Products)
Background:	Description: E3 ubiquitin ligase. Mediates the ubiquitination and subsequent proteasomal
	degradation of CKM, GMEB1 and HIBADH. Regulates the proteasomal degradation of muscle
	proteins under amino acid starvation, where muscle protein is catabolized to provide other
	organs with amino acids. Inhibits de novo skeletal muscle protein synthesis under amino acid
	starvation. Regulates proteasomal degradation of cardiac troponin I/TNNI3 and probably of
	other sarcomeric-associated proteins. May play a role in striated muscle atrophy and
	hypertrophy by regulating an anti-hypertrophic PKC-mediated signaling pathway. May regulate
	the organization of myofibrils through TTN in muscle cells.
	Gene: TRIM63
Molecular Weight:	40kDa
Gene ID:	84676
UniProt:	Q969Q1
Application Details	
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:100, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 $\%$
	glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months




Western Blotting

Image 1. Western blot analysis of extracts of mouse fetal muscle, using TRIM63antibody. The lane on the left is treated with the antigen-specific peptide.



Immunofluorescence (fixed cells)

Image 2. ABIN6277442 staining Hela cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600 was used as secondary antibod



Immunohistochemistry

Image 3. ABIN6277442 at 1/100 staining Mouse muscle tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22_{i} ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary