antibodies -online.com





anti-TAS2R10 antibody (Internal Region)



| () | 11/ | IN | /ie | A . |
|-----|-------------------|------|--------|--------|
| | / // | ۱ ات | / (− | ' \/\/ |
| | | | | |

| Quantity: | 50 μL |
|----------------------|---|
| Target: | TAS2R10 |
| Binding Specificity: | AA 100-180, Internal Region |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This TAS2R10 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunofluorescence (IF) |

Product Details

| Purpose: | Rabbit polyclonal to T2R10. |
|---------------|---|
| Immunogen: | Synthesized peptide derived from human T2R10 |
| Isotype: | lgG |
| Specificity: | T2R10 Polyclonal Antibody detects endogenous levels of T2R10 protein. |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |

Target Details

| Target: | TAS2R10 |
|-------------------|--------------------------|
| Alternative Name: | T2R10 (TAS2R10 Products) |

Target Details

| Molecular Weight: | 35 kDa |
|-------------------|--------|
| Gene ID: | 50839 |
| UniProt: | Q9NYW0 |

Application Details

| Application Notes: | WB 1:500-1:2000 |
|--------------------|---|
| | IF 1:200-1:1000 |
| | ELISA 1:10000 |
| Comment: | Expressed in subsets of taste receptor cells of the tongue and palate epithelium and exclusively in gustducin-positive cells. |
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 mg/mL |
| Buffer: | Liquid in PBS containing 50 % glycerol, 0.5 % BSA and 0.02 % sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C, and avoid repeat freeze-thaw cycles. |