antibodies -online.com





anti-SLC6A16 antibody (Internal Region)



Go to Product page

| () | 11/0 | K\ / | iew | 1 |
|-----|--------|------|---------------------|---|
| | \cup | ועוי | $\square \vee \vee$ | ı |
| | | | | |

| Quantity: | 50 μL |
|----------------------|--|
| Target: | SLC6A16 |
| Binding Specificity: | AA 210-290, Internal Region |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This SLC6A16 antibody is un-conjugated |
| Application: | ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

Product Details

| Purpose: | Rabbit polyclonal to NTT5. |
|---------------|---|
| Immunogen: | Synthesized peptide derived from human NTT5 |
| Isotype: | IgG |
| Specificity: | NTT5 Polyclonal Antibody detects endogenous levels of NTT5 protein. |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |

Target Details

| Target: | SLC6A16 |
|-------------------|-------------------------|
| Alternative Name: | NTT5 (SLC6A16 Products) |

Target Details

| Molecular Weight: | 82.2 kDa |
|-------------------|----------|
| Gene ID: | 28968 |
| UniProt: | Q9GZN6 |

Application Details

| Application Notes: | IHC 1:100-1:300 |
|--------------------|---|
| | ELISA 1:40000 |
| Comment: | Highly expressed in peripheral tissues, particularly in testis, pancreas, and prostate. |
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 mg/mL |
| Buffer: | Liquid in PBS containing 50 % glycerol, 0.5 % BSA and 0.02 % sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C, and avoid repeat freeze-thaw cycles. |